

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

10/590694

1

**AN ITRACONAZOLE EXHIBITING AN IMPROVED SOLUBILITY,
A METHOD OF PREPARING THE SAME AND A PHARMACEUTICAL
COMPOSITION FOR ORAL ADMINISTRATION COMPRISING THE SAME**

5

CROSS REFERENCE TO RELATED APPLICATION

This application is based on application No. 97-24938 filed in the Korean Industrial Property Office on June 16, 1997, the content of which is incorporated hereinto by reference.

10

BACKGROUND OF THE INVENTION

(a) Field of the Invention

The present invention relates to an itraconazole and, more particularly, to an itraconazole exhibiting an improved solubility, a method of preparing the same and a pharmaceutical composition comprising the same.

15 **(b) Description of the Related Art**

Because the water-solubility of an azole-type antifungal agent such as itraconazole is low, bioavailability of a drug including the azole-type antifungal agent is low, when the drug is orally administered to a patient. Various studies have been attempted to increase the water solubility and
20 bioavailability of the azole-type antifungal agent. The solubility and bioavailability of the compounds can be increased by complexation with cyclodextrins or derivatives thereof as described in WO 85/02767 and U. S. patent No. 4,764,604. WO 94/05263 reports a method of formulating a bead type drug by using a soluble polymer and the itraconazole.

As the azole-type antifungal agent, itraconazole has been widely used. Itraconazole has a molecular formula of $C_{35}H_{38}Cl_2N_8O_4$ and molecular weight of 705.64. Itraconazole is powder showing light yellow color and insoluble in water (solubility of less than 1 $\mu\text{g}/\text{mL}$), slightly soluble in alcohol (300 $\mu\text{g}/\text{mL}$) and soluble in methylene chloride (239 mg/mL). Itraconazole is a weak basic compound ($\text{pK}_a = 3.7$) and is only ionized at low pH, such as in gastric juice. The log partition coefficient of itraconazole in a system of n-octanol and an aqueous buffer solution of pH 8.1 is 5.66, indicative of very high lipophilicity. This property may influence its plasma protein binding and tissue distribution. Furthermore, it has been known that itraconazole is a broad spectrum antifungal compound developed for oral, parenteral and/or topical use and is disclosed in U. S. patent No. 4,267,179.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide itraconazole exhibiting an improved solubility by reducing particle size and changing the crystallinity thereof from crystalline into amorphous.

It is another object to provide a method of preparing the itraconazole by dissolution-induced drying.

It is another object to provide a pharmaceutical composition for oral administering the itraconazole.

These and other objects may be achieved by itraconazole exhibiting an improved solubility, wherein a particle diameter of the itraconazole is about 0.5 to about 10 μm and has an amorphous form.

The present invention further includes a method of preparing

itraconazole exhibiting the improved solubility. The method includes the steps of dissolving itraconazole in an organic solvent and dissolution-induced drying the mixture.

The present invention further includes a pharmaceutical composition including the itraconazole exhibiting the improved solubility as an active ingredient.

BRIEF DESCRIPTION OF THE DRAWINGS

A more complete appreciation of the invention, and many of the attendant advantages thereof, will be readily apparent as the same becomes better understood by reference to the following detailed description when considered in conjunction with the accompanying drawing, wherein:

FIG. 1 is a graph showing dissolution profile of itraconazole with and without dissolution-induced drying according to time; and

FIG. 2 is a graph showing X-ray diffraction patterns of itraconazole with and without dissolution-induced drying.

DETAILED DESCRIPTION OF THE INVENTION

For absorbing a solid drug into a systemic circulation via epithelial cells, the solid drug should be soluble in water. Because a water-insoluble drug is slowly dissolved from a solid formulation, the drug dissolution step is rate-limiting step in absorbing the drug. Thus, the potency of the drug and on set time, the potency duration time depend on the dissolution rate of the drug.

The concentration of a drug in blood depends on the absorption and

elimination rates thereof. Therefore, as the dissolution rate of the drug decreases, the absorption rate of the drug is reduced. As a result, although the total amount of drug absorbed into a human body is constant, time for reaching effective concentration thereof in blood is getting longer and maximum concentration thereof in blood decreases. Therefore, bioavailability and solubility of water-insoluble drugs decreases.

In order to solve these disadvantages and increase bioavailability and solubility, it is required to increase the dissolution rate of the water-insoluble drugs. For that purpose, particle size of the drug is reduced, or polymorphism, complex form, amorphous form, solid solution, co-melting mixture, conjugated compound, solvated compound of the drug and general drug additives are used.

The present invention provides an improved itraconazole having reduced particle size and modified crystalline characteristics from crystalline into amorphous. The improved itraconazole exhibits good water solubility, thereby increasing the bioavailability.

A method of preparing the improved itraconazole will be now illustrated in more detail.

Itraconazole is dissolved in an organic solvent and dissolution-induced dried to obtain powder of itraconazole. An amount of itraconazole in the organic solvent is 2 to 6 wt % and preferably, 4 wt %. It is preferred that the organic solvent is methylene chloride which is a good solvent for itraconazole. The dissolution-induced drying step is performed with a spray drier, a centrifugal granulator or fluid-bed granulator. In the dissolution-

induced drying step, a carrier, for example, lactose or other pharmaceutical excipients may be used. The pharmaceutical excipients may include a binder, a disintegrant, an agent for increasing viscosity, a lubricant, a stabilizer, a surfactant, a preservative, an electrolyte, a composite and a complex compound or another active material.

In the preparing step, it is important to control a dissolution-induced drying rate. If dissolution-induced drying rate is extremely slow, the solvent is too fast evaporated so that a large amount of drug is loss. Contrarily, if dissolution-induced drying rate is fast, microparticles are entangled with another, prior to evaporation, enlarging the particle size. It is preferred that the dissolution-induced drying rate is controlled to be slow initially and then controlled to gradually increase.

The particle size depends on an atomizing air pressure as well as the dissolution-induced drying rate. Low atomizing air pressure results in the formulation of larger droplets and increased tendency toward agglomeration. High atomizing air pressure could conceivably carry the risk of dissolution-induced drying of the drug solution, but this was found not to be a problem. Consequently, atomizing air pressure may be set at nearly maximum level.

The physical changes of itraconazole may be confirmed by differential scanning calorimeter (DSC) and x-ray crystallography.

Itraconazole prepared by method of the present invention has a particle diameter of about 0.5 to about 10 μm and an average particle diameter of about 3.7 μm . The itraconazole may include a trace of particles with diameter of below 0.5 μm .

The itraconazole of the present invention may be used as an active ingredient for a pharmaceutical composition for oral administration.

The pharmaceutical composition further includes vehicle such as lactose as carrier, or other pharmaceutical excipients.

-5 The pharmaceutical excipients may be include a binder, a disintegrant, an agent for increasing viscosity, a lubricant, a stabilizer, a surfactant, a preservative, an electrolyte, a composite and a complex compound or another active material.

10 Itraconazole of the present invention is applied to various types of a drug for oral administration. The various type of the drug are a tablet, a capsule, a granule and a fine-granule. One tablet or one capsule may include 50 mg to 100mg of itraconazole.

15 The present invention is further explained in more details with reference to the following examples. The examples is not intended to limit the present invention.

Example 1

12g of itraconazole was accurately weighed and dissolved in 200 ml of methylene chloride. The mixture was then dissolution-induced dried by using a spray-drier to obtain powder of itraconazole. The air temperature in
20 inlet and outlet of the spray drier were controlled to be $80 \pm 5^{\circ}\text{C}$ and $60 \pm 5^{\circ}\text{C}$, respectively. The dissolution-induced drying rate was controlled to be slow initially and then controlled to gradually increase.

Example 2

12g of itraconazole was accurately weighed and dissolved in 200 ml

of methylene chloride. The mixture was then dissolution-induced dried by using a fluid-bed granulator. The air temperatures in inlet and outlet of the fluid-bed granulator were controlled to be $80 \pm 5^\circ\text{C}$ and $60 \pm 5^\circ\text{C}$, respectively. The dissolution-induced drying rate was controlled to be slow initially and then controlled to gradually increase. During the dissolution-induced drying, a drying step was performed to completely evaporate the solvent.

Example 3

12g of Itraconazole was accurately weighed and dissolved in 200 ml of methylene chloride. The mixture was then dissolution-induced dried by using a C/F granulator to prepare powder thereof. The air temperatures in inlet and outlet of the C/F granulator were controlled to be $80 \pm 5^\circ\text{C}$ and $60 \pm 5^\circ\text{C}$, respectively. The rotation speed of rotor in the C/F granulator was 200 rpm. The dissolution-induced drying rate was controlled to be slow initially, and then controlled to gradually increase. During the dissolution-induced drying, a drying step was performed to completely evaporate the solvent.

Example 4: Tablet preparation

Itraconazole prepared from Example 1, lactose, calcium carboxymethylcellulose and hydroxypropylmethylcellulose were uniformly mixed. The mixture was dry-granulated by a roller compactor to prepare granules. The granules were mixed with magnesium stearate as lubricant. The mixture was tableted to prepare a tablet. The amounts of each of the components in one tablet were as follows.

	Itraconazole	100mg
	lactose	279mg
	calcium carboxymethylcellulose	250mg
	magnesium stearate	1mg
5	hydroxypropylmethylcellulose	20mg

Example 5: Capsule preparation

Itraconazole prepared from Example 1, lactose, calcium carboxymethylcellulose and hydroxymethylcellulose were uniformly mixed. The mixture was dry-granulated by using a roller compactor to prepare
10 granules. The granules were mixed with magnesium stearate as lubricant and charged into a hard capsule No. 0. The amounts of each of the components were as follows:

	Itraconazole	100mg
	lactose	179mg
15	calcium carboxymethylcellulose	200mg
	magnesium stearate	1mg
	hydroxypropylmethylcellulose	20mg

Experiment 1: Determination of solubility

Excess of itraconazole from Example 1 and conventional
20 itraconazole prepared without dissolution-induced drying were added to artificial gastric juice of pH 1.2 and shaken at 37°C for 2 hours, respectively. Each of the mixture were filtered through a membrane filter with pore diameter of 0.45 μ m and each of the filtrate was subjected to high-performance liquid chromatography, thereby obtaining gastric juice solubility.

The results are shown in Table 1.

Table 1: Solubility (37°C, 2hours, pH 1.2)

	$\mu\text{g/ml}$
The conventional itraconazole without dissolution-induced drying	3.5
Itraconazole of Example 1	218.6

As shown in Table 1, solubility of itraconazole of the Example 1 is 62
5 times larger than that without dissolution-induced drying. Therefore, it is shown that solubility of itraconazole of the present invention significantly increases with compared to the conventional itraconazole without dissolution-induced drying.

Experiment 2: Determination of particle size

10 Itraconazole powder prepared from Example 1 and the conventional itraconazole powder without dissolution-induced drying were uniformly distributed in liquid paraffin and particle diameter thereof were determined, respectively. The results are shown in Tables 2 to 4.

15 Table 2: The conventional Itraconazole without dissolution-induced drying

Particle size (μm)	Distribution ratio of particle (%)	Accumulated amount (wt %)
~5	2.4	2.4
5~15	21.8	24.2

15 ~ 17.5	18.4	40.6
17.5 ~ 20	9.9	50.5
20 ~ 30	23.9	74.4
30 ~ 35	8.0	82.4
35 ~ 40	8.1	90.5
40 ~ 52.5	9.5	100.0

Table 3: Itraconazole of Example 1

Particle size (μm)	Distribution ratio of particle (%)	Accumulated amount (wt %)
~0.5	2.4	2.4
0.5 ~ 1.25	25.8	28.2
1.25 ~ 2.5	24.0	52.2
2.5 ~ 5	34.2	86.4
5 ~ 7.5	12.2	98.6
7.5 ~ 10	1.4	100.0

Table 4: Average particle diameter

	μm
The conventional itraconazole without dissolution-induced drying	24.5
itraconazole of Example 1	3.7

The average diameter of itraconazole particle of the present invention is 7 times smaller than that without dissolution-induced drying .

Experiment 3: Dissolution rate test

The dissolution rate test on the itraconazole tablet prepared from Example 4 and a conventional itraconazole tablet without dissolution-induced drying were performed.

The itraconazole tablet prepared from Example 4 and the conventional itraconazole tablet were respectively added to artificial gastric juice of pH 1.2 as an eluting solvent. While keeping at 37°C, 5ml of the sample was taken from the mixture after 5, 10, 30, 60 and 120 minutes, respectively. Each of the samples were filtered by using a filter of 0.45 micrometers and each of the filtrate were subjected to high-performance liquid chromatography. The results are shown in Fig. 1.

As shown in Fig.1, dissolution rate and solubility of itraconazole with dissolution-induced drying is greatly enhanced with compared to itraconazole without dissolution-induced drying.

Experiment 4: The physical property test

The physical changes of itraconazole was confirmed by x-ray diffraction and result are shown in Fig. 2. In Fig. 2, pattern A shows x-ray diffraction pattern of itraconazole without dissolution-induced drying. Pattern B shows x-ray diffraction pattern of itraconazole with dissolution-induced drying. As shown in Fig. 2, itraconazole without dissolution-induced drying has an crystalline form and itraconazole with dissolution-induced drying has an crystalline form.

As described above, the present invention is capable of increasing solubility of itracobazole being insoluble drug and a dissolution rate and itraconazole of the present invention is applied to solid preparation.


While the present invention has been described in detail with
5 reference to the preferred embodiments, those skilled in the art will appreciate that various modifications and substitutions can be made thereto without departing from the spirit and scope of the present invention as set forth in the appended claims.

WHAT IS CLAIMED IS:

1. An itraconazole exhibiting an improved bioavailability wherein the particle diameter of itraconazole is in the range of 0.5 to 10 μm and the itraconazole has an amorphous form.
- 5 2. The itraconazole of claim 1 wherein an average particle diameter of itraconazole is about 3.7 μm .
3. A method of preparing an improved itraconazole having particle diameter of 0.5 to 10 μm and an amorphous form, comprising the steps of:
 - 10 dissolving an itraconazole in an organic solvent;
dissolution-induced drying the mixture.
4. The method of claim 3 wherein an average particle diameter of itraconazole is about 3.7 μm ,
5. The method of claim 3 wherein the dissolution-induced
15 drying step is performed by using a machine selected from the group consisting of spray drier, fluid-bed granulator and centrifugal granulator.
6. A pharmaceutical composition for oral administration, comprising an improved itraconazole having a particle diameter of 0.5 to 10 μm and having an amorphous form, as active ingredient.
- 20 7. The pharmaceutical composition of claim 6 wherein an average particle diameter of itraconazole is about 3.7 μm .
8. The pharmaceutical composition of claim 6 further comprising vehicle or pharmaceutical excipients.


1/2

FIG. 1



2/2

FIG.2



5 10 25 35 (2θ)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 98/00164

A. CLASSIFICATION OF SUBJECT MATTER

IPC⁶: C 07 D 403/14; A 61 K 31/41

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC⁶: C 07 D 403/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

AT, Chemical Abstracts

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO: WPI; Questel: CAS, Medline

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 147 171 A1 (TOKYO TANABE CO.) 03 July 1987 (03.07.87), claims; examples 1,2.	1-8
Y	WO 94/16 700 A1 (SEPRACOR INC.) 04 August 1994 (04.08.94), page 9, lines 13-25.	1-8
Y	EP 0 006 711 A1 (JANSSEN PHARM.) 09 January 1980 (09.01.80), claim 1.	1-8
A	EP 0 197 571 A1 (JANSSEN PHARM.) 15 October 1986 (15.10.86).	1-8
A	WO 94/05 263 A1 (JANSSEN PHARM.) 17 March 1994 (17.03.94).	1-8
A	WO 94/16 733 A1 (CHIESI FORM.) 04 August 1994 (04.08.94).	1-8
A	WO 95/31 178 A1 (JANSSEN PHARM.) 23 November 1995 (23.11.95).	1-8
A	WO 97/18 839 A1 (JANSSEN PHARM.) 29 May 1997 (29.05.97).	1-8

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

18 August 1998 (18.08.98)

Date of mailing of the international search report

26 August 1998 (26.08.98)

Name and mailing address of the ISA/
Austrian Patent Office
Kohlmarkt 8-10; A-1014 Vienna
Facsimile No. 1/53424/535

Authorized officer

Hammer

Telephone No. 1/53424/374

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 98/00164

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>Chemical Abstracts, Vol.125, No.26, 23 December 1996 (Columbus, Ohio, USA), page 708, column 1, abstract No.338841t, deCHASTEIGNER, S. et al.: "Comparative study of the association of itraconazole with colloidal drug carriers", & Drug Dev. Res. 1996, 38(2), 125-133.</p> <p style="text-align: center;">----</p>	1-8

Form PCT/ISA/210 (continuation of second sheet) (July 1993)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR 98/00164

Im Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
EP A1 147171		AU A1 36537/84 EP A2 147171 EP A3 147171 JP A2 60139688	04-07-85 03-07-85 28-01-87 24-07-85
WD A1 9416700	04-08-94	AU A1 60960/94 AU A1 59665/98 CA A1 2154777 EP A1 680324 EP A4 680324 JP T2 8509464 US A 5474997	15-08-94 28-05-98 04-08-94 08-11-95 10-04-96 08-10-96 12-12-96
EP A1 6711	09-01-80	US A 4267179 AT E 5140 AU A1 48160/79 AU B2 528095 BG A2 50387 BG B2 60430 CA A1 1149386 CY A 1251 DE CO 2966346 DK A 2614/79 DK E 160428 DK C 160428 EP B1 6711 ES A1 481771 ES A5 481771 FI A 791989 FI B 65429 FI C 65429 GR A 64910 HK A 749/84 HR B1 930473 HU B 187744 IE B 48762 IL A0 57625 IL A1 57625 JP A2 55011578 JP B4 63053192 JP A2 1211561 JP B4 5009431 ND A 792111 NO E 153297 NO C 153297 NZ A 190746 PH A 16198 PT A 69805 RD P 77325 SG A 451/84 SI A8 7911479 YU A 1478/79 YU B 42188 US A 4368200 US A 4490530 CS A2 7904087 CS B2 241468 PL A1 2165553 PL B1 117988 SU A3 1069625 ZA A 7903128	12-05-81 15-11-84 03-01-80 14-04-80 15-07-80 31-03-80 05-07-80 31-08-80 01-12-80 24-12-79 11-04-81 19-08-81 26-10-82 01-07-80 01-08-80 24-10-79 31-01-84 10-05-84 09-06-80 12-10-84 31-10-86 28-02-86 15-05-86 31-10-79 30-12-80 26-01-80 21-10-88 24-08-80 04-02-80 28-12-79 11-11-80 19-02-86 15-12-80 28-07-80 01-06-80 17-08-81 08-03-85 31-10-80 31-12-80 30-06-80 11-01-84 25-12-80 15-08-80 13-04-80 24-04-80 30-09-81 24-01-84 25-02-81
EP A1 197571		AT E 124055 AT E 159954 AU A1 54760/86 AU B2 583599 CA A1 1300130 CY A 1918 DE CO 3650336 DE T2 3650336 DE CO 3650655 DE T2 3650655 DK A0 1214/86 DK A 1214/86 EP A2 197571 EP A5 197571 EP A1 636634 EP B1 197571 EP B1 636634 EP A1 552984	15-07-85 15-11-87 16-10-86 04-05-86 05-05-87 07-03-87 27-07-87 02-11-87 11-12-87 12-03-88 14-03-86 16-09-86 15-10-86 30-09-87 01-02-88 21-06-87 05-11-87 16-05-87

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR 98/00164

ES	A5	552984	15-06-87
ES	A1	8705899	01-08-87
FI	A0	861064	14-03-86
FI	A	861064	14-03-86
FI	A0	894274	11-09-89
FI	B	84076	28-06-84
FI	B	84076	10-10-84
FI	B	86507	29-05-86
FI	B	86507	10-09-86
GB	A0	8506792	17-04-85
GR	A	860685	11-07-86
HK	A	1318796	26-07-86
HU	A2	42514	28-07-87
HU	B	198092	28-07-87
HU	B	206381	28-10-87
IE	B	66673	24-01-86
JP	A2	61275301	03-12-81
JP	B2	9307386	09-11-93
KR	B1	860971	16-09-86
NO	A	167578	13-08-86
NO	B	167578	20-11-86
NZ	A	215350	29-03-89
PT	A	82186	01-04-82
PT	B	82186	21-04-82
US	A	4764604	16-08-86
US	B1	4764604	13-06-86
ZA	A	8601930	28-10-86
US	A	4870060	26-09-89
<hr/>			
WO A1	9405263	17-03-94	
AP	A0	9300563	31-10-93
AP	A	145444	19-01-94
AT	E	145444	19-01-94
AU	B2	665867	18-01-86
CN	A	1088432	29-06-89
CN	B	1034714	10-04-89
CN	A	9500543	14-08-95
DE	C0	69306119	03-01-89
DE	T2	69306119	12-01-89
DK	T3	6581033	09-12-85
EP	A1	6581033	21-06-85
EP	B1	6581033	21-11-85
ES	T3	2097536	01-04-86
FI	A	950975	02-03-95
FI	A0	950975	02-03-95
GR	T3	3032198	11-03-89
HR	A1	931158	20-06-93
HU	A0	9500642	28-04-95
HU	A2	70419	28-04-95
IL	A0	106871	28-10-86
IL	A1	106871	28-10-86
JP	T3	93051093	06-03-93
MX	A1	93051093	06-03-93
NO	A0	9500642	03-01-95
NO	A	9500642	03-01-95
NZ	A	9500642	03-01-95
PL	A1	172677	28-06-86
PL	B1	172677	28-06-86
PL	B1	174178	28-06-86
SI	A	93006461	11-03-93
US	A	9500642	27-03-95
ZA	A	9500642	02-03-95
<hr/>			
WO A1	9531178	23-11-95	
AP	A0	9600873	31-10-96
AU	A1	25226479	05-11-96
AU	B2	9580767	07-08-95
BR	A	95007873	07-08-95
CA	A	2118986	11-01-96
CN	A	96007873	07-08-95
CN	A	96007873	07-08-95
EP	A1	95007873	07-08-95
FI	A	964589	11-01-96
FI	A0	964589	11-01-96
HU	A0	9603188	28-08-96
HU	A2	763226	11-08-96
IL	A0	1137422	01-08-96
JP	T2	10300101	06-01-96
NO	A0	9648533	11-01-96
NO	A	9648533	11-01-96
NZ	A	9648533	11-01-96
PL	A1	285771	01-08-86
SK	A3	1469796	07-08-97
<hr/>			
WO A1	9718839	29-05-97	
AU	A1	76943796	11-06-97
NO	A0	981078	11-03-98

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)